



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07F 15/00, C07H 21/00, 23/00	A1	(11) International Publication Number: WO 98/15564 (43) International Publication Date: 16 April 1998 (16.04.98)
(21) International Application Number: PCT/NL97/00559 (22) International Filing Date: 8 October 1997 (08.10.97) (30) Priority Data: 96202792.6 8 October 1996 (08.10.96) EP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant (for all designated States except US): KREAT-ECH BIOTECHNOLOGY B.V. [NL/NL]; Keienbergweg 3, NL-1101 EZ Amsterdam (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): HOUTHOFF, Hendrik, Jan [NL/NL]; Schellingwouderdijk 125, NL-1023 NB Amsterdam (NL). REEDIJK, Jan [NL/NL]; Anthoni Duycklaan 4, NL-2334 CD Leiden (NL). JELSMA, Tinka [NL/NL]; Kerseboomstraat 30, NL-1326 DG Almere (NL). HEETE-BRIJ, Robert, Jochem [NL/NL]; Zadelstraat 2, NL-3511 LT Utrecht (NL). VOLKERS, Herman, Hendricus [NL/NL]; Noordervesting 3, NL-1141 SL Monnickendam (NL). (74) Agent: SMULDERS, Th., A., H., J.; Vereenigde Octrooi-bureaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHODS FOR LABELING NUCLEOTIDES, LABELED NUCLEOTIDES AND USEFUL INTERMEDIATES		
(57) Abstract <p>The present invention provides improved methods for labeling nucleotides. The method of the invention for labeling nucleotides comprises the steps of: reacting a reactive moiety of a linker, which linker is a platinum compound having a stabilizing bridge and two reactive moieties, with an electron donating moiety of a spacer, which spacer comprises a chain having at least four atoms and at least one heteroatom in the chain, which spacer further comprises said electron donating moiety at one end of the chain and a reactive moiety at the other end of the chain; reacting the reactive moiety of said spacer with a label; reacting the other reactive moiety of said linker with a nucleotide. A major advantage of the invention is that all nucleotides can be labeled by the method of the invention, whereas until now the attachment of a label was mostly restricted to one or certain nucleotides.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Title: Methods for labeling nucleotides, labeled
 nucleotides and useful intermediates.

The invention relates to methods for labeling nucleotides using linkers (linking moieties between labels and bio-organic molecules, which linkers are based on platinum compounds).

Platinum (coordination) compounds have been considered
5 interesting molecules for a very long time. For a review of
these compounds and their uses we refer to Reedijk et al.
(Structure and Bonding 67, p.53-89, 1987). Especially Cis-
platinum has received a lot of attention as a possible anti-
tumour drug. This anti-tumour reactivity of platinum compounds
10 originates from their having at least two reactive groups
(preferably cis-oriented towards each other), which make it
possible to cross-link DNA molecules, thereby inhibiting the
replication of these DNA molecules.

The British patent application 2 148 891 discloses cis-
15 platinum complexes, which are six-coordinated. The platinum is
attached to two halogens or hydroxy groups, two additional
halogens and to an ethylene diamine derived group, such as
1,2-diamino-2-methylpropane or 1,2-diamino-2-methylbutane. The
complexes are said to have an excellent anti-tumor effect.

20 In the European patent application four-coordinated
complexes of platinum to 2,3-alkyl-1,4-butanediamine and two
halogens are described for their anti-tumor effect.

Different four-coordinated platinum complexes are
described in the European patent application 0 386 243. The
25 complexes comprise a diamine bidentate ligand and two 2-
arylalkanoic acid or 3-aryl-2-oxoalkanoic acid ligands. These
complexes are stated to have a strong growth inhibiting action
on certain leukemia cells and are used for their oncostatic
activity.

30 US patent 4,207,416 discloses ethylenediamine-
platinum(II) 2,4-dioxypyrimidine complexes as having a high
anti-tumor activity and low mammalian toxicity.

A different use of platinum (coordination) compounds has been disclosed in PCT application (WO92/01699) wherein a platinum compound having only two reactive moieties (denominated as leaving groups therein) is reacted with a
5 fluorescein to obtain a labeled platinum compound which can bind (non-covalently) to a nucleic acid, preferably at the N-7 position of a guanine residue.

Several methods for labeling nucleotides have been described in the literature. For a long time, the standard
10 method has been to use radioactive isotope labeling. However, there are a number of problems associated with the use of radioisotopes, such as potential health hazards, disposal problems and instability problems.


In order to overcome these problems, Dale et al.,
15 Biochemistry, 14, (1975), 2447-2457, have proposed to use direct covalent mercuration as a labeling technique for nucleotides and polynucleotides. It was found, that cytosine and uracil may be mercured at their C5-position under mild conditions. Further, Gebeyehu et al., Nucleic Acids Research,
20 15, (1987), 4513-4534, have reported that adenine and cytosine may be labeled with biotin derivatives through an aliphatic linker of from 3 to 17 atoms.

A major drawback of these known methods is that they are not suitable for labeling all different nucleotides. For
25 instance, Dale et al. reported that their covalent mercuration method leads to negative results for adenine, thymine and guanine bases. In some cases, for example when only a few residues of a certain nucleotide are present in a certain nucleic acid or when the terminating nucleotide residue of a
30 nucleic acid has to be labeled, it is desired to have at one's disposal a method for labeling any nucleotide residue.

The present invention provides such a method. The method for labeling nucleotides of the invention comprises the steps of:

35 - reacting a reactive moiety of a linker of the formula

3



- wherein X represents any stabilizing bridge and wherein A and B represent the same or different reactive moieties, with an electron donating moiety of a spacer, which spacer comprises a chain having at least four atoms and at least one heteroatom in the chain, which spacer further comprises said electron donating moiety at one end of the chain and a reactive moiety at the other end of the chain;
- 10 - reacting the reactive moiety of said spacer with a label;
 - reacting the other reactive moiety of said linker with a nucleotide.

According to the invention, the linker may first be attached to the nucleotide and then to the spacer, or vice versa and the spacer may first be coupled to the label and then to the linker or vice versa.

The reactive moiety of the spacer may be any reactive moiety that will enable the reaction between the spacer and the label in such a manner that a labeling moiety comprising a label and a spacer is formed, which labeling moiety is sufficiently stable.

The main purpose for labeling nucleotides is that these labeled nucleotides can be incorporated in nucleic acid molecules. Modified nucleotides, especially those wherein a (bulky) label is attached to the nucleotide, are often built-in into nucleic acids with a much lower efficiency. The methods according to the invention result in labeled nucleotides which are built-in into nucleic acids with a higher efficiency than the labeled nucleotides available to date. This is probably for due to the selection of the spacers according to the invention in combination with the platinum-based linkers according to the invention.

The label to be used according to the invention is not critical. In principle all labels which can be attached to a nucleotide and are employed to date can be used. These labels may be radioactive labels, enzymes (which need reaction with a substrate to be detected), specific binding pairs components such as avidin, streptavidin or biotin, biocytin, iminobiotin, colloidal dye substances, fluorochromes (rhodamin, etc.), reducing substances (eosin, erythrosin, etc.), (coloured) latex sols, digoxigenin, metals (ruthenium), metal sols or other particulate sols (selenium, carbon and the like), dansyl lysin, Infra Red Dyes, coumarines (amino methyl coumarine), antibodies, protein A, protein G, etc. The invention has most benefits with bulkier labels such as biotin, avidin, streptavidin, digoxigenin or a functional equivalent thereof.

The invention is not limited to nucleotides or nucleosides as such; derivatives and functional equivalents are also included. The usual nucleotides adenine, thymidine, cytosine, guanine and uridine are preferred. Especially the purines are preferred which have a very good incorporation rate.

For coupling of the spacer to the platinum linker an electron donating moiety is required. In a preferred method the electron donating moiety is an amine or a thiolate anion, which have both proven to be very successful. It was found that aromatic amines, such as imidazoles or purines, are capable of forming very strong bonds to platinum and thus are very suitable for use as the electron donating moiety.

The spacer is an important aspect of the present invention; it provides the easiest coupling between label and linker. For avoiding steric hindrance in incorporation of the nucleotide into the nucleic acid it should at least be four atoms long, preferably it is at least four carbon atoms long and has at least one heteroatom in that carbon chain. A heteroatom confers a certain amount of rigidity on the spacer. This rigidity provides an additional assurance that steric factors will not obstruct a convenient linking of a nucleotide and a label. It is preferred that at least one heteroatom is


an oxygen atom, which positively effects the hydrophilicity of the spacer.

Preferably, the spacer comprises no more than 20 carbon atoms in the chain, which is preferably an essentially non-branched chain, thus causing no steric hindrance. The reason for this will be clear.

A highly preferred spacer is 1,8-diamino-3,6-dioxaoctane, herein referred to as Dadoo. Dadoo is a very flexible compound with a distal primary amine group and a size that makes it very suitable for use as spacer according to the invention.


Another highly preferred spacer of the invention is an oligolysine or a polylysine. Due to their structure and conformation, these molecules create the most convenient environment for an optimal interaction among the actual label, the nucleotide and the platinum. An additional advantage of the use of lysine chains as the spacer is, that by altering the number of lysine units in the chain, the optimal conditions for specific labels and nucleotides or nucleic acids can be attained. Given a certain application, the skilled person will easily determine how many lysine units are required for optimum results.

An especially interesting labeling moiety comprising a label and a spacer, has the formula




25

or the formula




wherein X represents any stabilizing bridge, Z and Z' represent a non-leaving ligand and n is an integer of from 2 to 10.

Accordingly, the linker-spacer-label-system, or labeling substance, with the labeling moiety of formula (II) or formula (III) has the formula



10

or the formula




wherein A, X, Z, Z' and n have the above meanings.

The non-leaving ligands Z and/or Z' are preferably an
 5 NH₃, NH₂R, NHR₂ or NR₃ group, wherein R represents an alkyl
 group having from 1 to 6 carbon atoms, because these ligands
 have an even smaller leaving-group character than other non-
 leaving ligands.

The interesting feature of using the labeling moieties
 10 having formulas (II) and (III) is that both the nucleotide and
 the actual label have the benefit of being bonded directly to
 a platinum atom, while at the same time these moieties are
 sufficiently far apart to avoid steric hindrance.

The linkers according to the invention preferably are
 15 platinum compounds wherein X represents an aliphatic diamine.
 In a preferred embodiment of the invention, one or both of the
 nitrogen atoms of the aliphatic diamine are shielded. A
 suitable manner of shielding these nitrogen atoms consists of
 substitution with one or two alkyl groups of from 1 to 6
 20 carbon atoms, preferably methyl groups. This is advantageous
 in that hydrogen bonding between the triphosphate group of a
 nucleotide and the stabilizing bridge is prevented.
 Preferably, a diamine having 2-6 carbon atoms is used, most
 preferably an ethylene diamine group, which is well-known for

its stabilizing effect on this class of platinum compounds. In this case, the linker has the formula




5

wherein G represents hydrogen or an alkyl group of from 1 to 6 carbon atoms and A and B represent the same or different reactive moieties.

10 The coupling or reactive moieties A and B are preferably the same and selected from the group consisting of NO_3^- , SO_3^- , Cl^- , I^- , or other halogens.

The invention of course also encompasses a labeled nucleotide obtainable by a method as disclosed above.

15 In addition, the invention encompasses a labeling substance for labeling nucleotides by a method as disclosed above. The labeling substance of the invention has the formula



20 wherein X and A have the above meanings and the labeling moiety comprises a label and a spacer as described above. Of course the labeling substances of the invention can also be used for labeling purposes other than labeling nucleotides. It was found that numerous (bio-) organic compounds, i.e. nearly
 25 every bio-organic molecule which contains an accessible

sulphur or nitrogen atom, for example proteins, can be labeled with the platinum compounds of the invention.

A great advantage of the invention arises from the use of the platinum compounds having formula (I) as linkers in the methods of preparing labeled nucleotides according to the invention. These linkers can be prepared by very convenient and reliable methods.

In WO92/01699 the starting compounds disclosed for preparing labeled platinum compounds are platinum(II)(ethylenediamine)dichloride and platinum(II)(ethylenediamine)(Me₂SO)Cl. The first one can be obtained commercially, the second one (the preferred one) must be synthesized. In the dichloride compound, the Cl⁻ions are less readily substituted by a label or a nucleotide, respectively. In the latter case, the total nucleotide labeling time will be appreciably longer, up to several hours, instead of several minutes.

The methods for preparing the linkers that are used in the method of labeling nucleotides according to the invention are based on the selection of suitable starting compounds of the formula PtE₄ wherein E is an electronegative group, preferably a halogen or NO₃⁻ or SO₃⁻. The reaction, which is described in the examples, of these starting compounds with e.g. ethylenediamine is very simple and efficient. Moreover, this reaction leads to very suitable symmetric intermediate compounds for producing labeled nucleotides. A major advantage of using these compounds is that when a stabilizing bridge for the resulting platinum compound has to be attached that no blocking reagents have to be employed. Another advantage is that the resulting intermediate compounds can again be labeled without the use of blocking agents. Therefore steps removing blocking agents can be eliminated completely. Furthermore the yields of these reactions are very high. Yet another advantage of the use of these symmetrical starting compounds is that no mixtures of different resulting compounds can be formed, which may interfere with the following reaction and reduce yield or require extra separation steps.

A very suitable intermediate compound according to the invention is platinum(II)(ethylenediamine)(NO₃)₂. This substance can very easily be provided with a suitable spacer and a labeling group, resulting in labeling substances which
5 can, through substitution of the remaining NO₃-group be linked to a nucleotide quite easily. Furthermore the methods for producing these compounds and the resulting compounds do not involve highly toxic substances such as DMSO.

The intermediate compounds can be labeled with any
10 suitable label (also known as marker) through a spacer as disclosed hereinabove.

Furthermore, the known advantages (from WO92/01699 for instance) are of course also obtained with the present methods and compounds. Another advantage of the platinum compounds is
15 that they can be detected more or less directly by using the platinum as a nucleus for depositing silver or other metal crystals.

By binding the labeling substance to a nucleotide residue, DNA or RNA molecules, be it single stranded or
20 otherwise, can be easily detected, but it also allows for the production of probes for hybridization techniques wherein unlabeled DNA/RNA molecules hybridize to the labeled probe. The platinum linker labeled nucleotides do hardly interfere with the hybridization, if at all. Also, this technique
25 obviates the use of modified nucleotides in preparing probes.

Nucleotides modified in accordance with the practices of this invention and oligo- and polynucleotides into which the modified nucleotides have been incorporated or oligo- and
30 polynucleotides that have been directly modified using these novel platinum compounds may be used as probes in biomedical research, clinical diagnostics and recombinant DNA technology.


Other advantages and embodiments of the invention will become clear from the following experimental part and the examples.

EXPERIMENTAL

SYNTHESIS OF INTERMEDIATE PLATINUM COMPOUNDS


5 These compounds, i.e. the linkers having formula (I), may be prepared by a process which involves:

(a) reacting a compound having the structure:



10


with KI in a suitable solvent under suitable conditions so as to form a iodated platinum compound having the structure:




15

(b) reacting said iodated platinum compound with ethylenediamine in a suitable solvent so as to form a diethylenediamine iodated platinum compound and represented by the formula $\text{Pt}(\text{en})\text{I}_2$ and having the structure:

20




(c) reacting said compound with AgNO_3 , the reaction being carried out in a suitable solvent, under suitable conditions so as to form a compound having the structure:




5

(d) reacting said compound with KCl in a suitable solvent under suitable conditions so as to form a compound having the structure:



10

(e) reacting said compound with AgNO_3 in a suitable solvent, under suitable conditions so as to form a compound having the structure:



15

(f) recovering said compound as modified platinum starting compound for the synthesis of hapten-bound $\text{Pt}(\text{en})$ compounds for use as DNA and/or RNA label.

20

Example 1

A. Preparation of Pt(en)-diamine starting material.

- 5 Preparation of Platinumethylenediamine(NO₃)₂: starting material.
Pt(en)(NO₃)₂

All reactions are performed in the dark.

- 10 Dissolve 1 gram potassium tetrachloroplatinate (II), K₂PtCl₄ (2.4 mmol, Sigma) in 50 ml millipore (filtered water) and stir at room temperature. Add 10 equivalents of potassium iodide, KI (24 mmol, 3.999 g, Sigma). The colour of the solution will immediately turn from orange into dark red
15 (K₂PtI₄), stir for 5 minutes.

Add one equivalent ethylenediamine (2.4 mmol, 160.8743 µl, Merck 11=0.9 kg) after diluting 161 µl ethylenediamine in 5 ml millipore very slowly to the platinum solution, mix this solution for 1 hour at room temperature.

- 20 A yellow/brown precipitate, Pt(en)I₂, will be formed and the liquid standing above should be clear.

- Filter the solution through a 1.0 µm membrane filter (Schleicher&Schuell), wash the precipitate with millipore, ethanol and diether (in this order). Dry the Pt(en)I₂ for at
25 least 4 hours in a vacuum dryoven at 37°C.

Weigh the dried Pt(en)I₂ (~1.07 g) and suspend it in 45 ml millipore/5 ml acetone, the solution will be cloudy. Add 1.95 equivalent of AgNO₃ (M = 169.9, Sigma). Stir the reaction overnight at room temperature.

- 30 Filter the solution through a 1.0 µm membrane filter, the precipitate is Silveriodide, AgI, the filtrate should be clear.

- Add to 0.5 ml of the filtrate, Pt(en)(NO₃)₂, an excess of KCl or NaCl and make sure that no white precipitate is formed
35 immediately after adding the excess of KCl or NaCl. If no white precipitate (only a yellow one) is formed then add an excess of KCl or NaCl to the entire filtrate. After the yellow precipitate is formed filter the solution and wash the precipitate (Pt(en)Cl₂) with millipore, ethanol and diether.

Dry the precipitate for at least 4 hours in a vacuum dryoven at 37°C.

Weigh the dry Pt(en)Cl_2 ($M=326,1$), and suspend it in 45 ml millipore/5 ml aceton and stir the cloudy suspension. Add 1.95
5 equivalent AgNO_3 and stir the solution overnight at room temperature. The colour of the solution will become white, due to the formation of AgCl .

Filter the solution in the dark and evaporate the filtrate to remove the aceton by rotation evaporation untill
10 25 ml of the filtrate is left. The filtrate is then freeze-dried. The product is checked by NMR or Infrared Absorption Spectroscopy.

B. Preparation of Pt(tmen) -diamine starting material.

15

Preparation of Platinum- N,N,N',N' -tetramethylethylenediamine(NO_3)₂: starting material.
 $\text{Pt(tmen)}(\text{NO}_3)_2$

20 All reactions are performed in the dark

Repeat the entire procedure of Example 1A, but use N,N,N',N' -tetramethylethylenediamine instead of ethylenediamine.

Example 2

25

A. Preparation of $[\text{Pt(en)}(\text{BioDadoo-NH}_2)(\text{NO}_3)](\text{NO}_3)$

Dissolve $\text{Pt(en)}(\text{NO}_3)_2$ (18.2 mg, 0.048 mmol) in 10 ml of Millipore water and heat until dissolving. Dissolve BioDadoo
30 (20 mg, 0.053 mmol, purchased from Boehringer Mannheim) in 5 ml of Millipore water. Add the two solutions together and adjust the pH to 8 by 0.1 N NaOH, react for at least 3 hours at 50°C. Isolate the end product by freeze drying.

B. Preparation of $[Pt(tmen)(BioDadoo-NH_2)(NO_3)](NO_3)$

Dissolve $Pt(tmen)(NO_3)_2$ (35 mg, 0.08 mmol) in 12.5 ml of Millipore water and heat until dissolving. Dissolve BioDadoo (32 mg, 0.085 mmol) in 10 ml of Millipore water. Add the two solutions together and adjust the pH by 0.1 N NaOH, react for at least 4 hours at 50°C. Isolate the end product by freeze drying.

10 C. Preparation of $[Pt(en)(DigDadoo-NH_2)(NO_3)](NO_3)$

Dissolve $Pt(en)(NO_3)_2$ (5 mg, 0.013 mmol) in 5 ml of Millipore water and heat until dissolving. Dissolve DigDadoo (9 mg, 0.016 mmol, purchased from Boehringer Mannheim) in 5 ml of Millipore water. Add the two solutions together and react for at least 4 hours at 50°C. Isolate the endproduct by freeze drying.

Example 3

20

A. Preparation of a labeled dGTP

Dissolve $[Pt(en)(BioDadoo-NH_2)(NO_3)](NO_3)$ (9 mg, 0.012 mmol) in 2 ml of Millipore water. Add 2'-deoxyguanosine-5'-triphosphate (2.3 mg, 0.004 mmol) and adjust the pH to 6. Incubate for 24 hours at 50°C, freeze-dry and redissolve in Millipore water (1 ml) and filter through a membrane filter. Apply the mixture to a FPLC with MonoQ and purify with a linear gradient from 100% Millipore water to 100% 1M NH_4HCO_3 , collect and pool appropriate fraction and isolate by freeze drying. Dissolve the product in a 100 mM solution of triethylamine ammonium acetate (TEAA) (1 ml) and apply to a Reversed Phase HPLC (C18) with a linear gradient from 100% 100 mM TEAA to 50% 100 mM TEAA / 50% 100 mM TEAA/acetonitrile (1/1 v/v), collect and pool the appropriate fraction and remove solvents by repeated evaporation in vacuo. Pass the product over a cation exchanger

(Dowex) in the lithium form, isolate the product by freeze drying.

B. Preparation of a labeled 5-AA-dUTP

5

Dissolve [Pt(en)(BioDadoo-NH₂)(NO₃)](NO₃) (6 mg, 0.008 mmol) in 2 ml of Millipore water. Add 2'-deoxyuridine-5-aminoallyl-5'-triphosphate (2 mg, 0.004 mmol) and adjust the pH to 8.

- Incubate for 24 hours at 50°C, freeze-dry and redissolve in
10 Millipore water (1 ml) and filter through a membrane filter.
Apply the mixture to a FPLC with MonoQ and purify with a linear gradient from 100% Millipore water to 100% 1M NH₄HCO₃, collect and pool appropriate fraction and isolate by freeze drying. Dissolve the product in a 100 mM solution of
15 triethylamine ammonium acetate (TEAA) (1 ml) and apply to a Reversed Phase HPLC (C18) with a linear gradient from 100% 100 mM TEAA to 50% 100 mM TEAA / 50% 100 mM TEAA/acetonitrile (1/1 v/v), collect and pool the appropriate fraction and remove solvents by repeated evaporation in vacuo. Pass the product
20 over a cation exchanger (Dowex) in the lithium form, isolate the product by freeze drying.

Example 4

25 Reaction for coupling Pt(en)-compounds to DNA

Typical reaction for labeling DNA molecules with a Pt-compound according to the invention.

- 30 5 µg of double stranded DNA is sonicated or DNase treated to yield fragments of 300-500 bp. 6 µg of Pt(en)-compound is added and the volume is adjusted to 50 µl with demineralised water. The reaction mixture is incubated at 65°C for 1 hour. Non-bound Pt(en)-compound is blocked by adding 100 µl of a
35 NADDTc solution. The Pt(en)-compound labeled DNA is purified on a sphadex G-50 column. Readily labeled and purified DNA is stored at -20°C or used directly in a DNA probe based assay.

Pt(en)-compound labeled DNA probes can be stored at least 2 years at -20°C without loss of activity and/or specificity. All applications mentioned are carried out with probes labeled according to this protocol.

5

Example 5

Biotin labeling of DNA probes with [Pt(en)(BioDadoo-NH₂)(NO₃)](NO₃) (BioDadoo-ULS).

10

Introduction

The labeling method has been used to label DNA probes with Biotin for In Situ Hybridization (ISH). In this example labeling procedures including the protocols and data for quality control procedures are presented. For Biotin labeling a plasmid cloned total DNA of Human Papilloma Virus type 6 (HPV-6, 40% GC basepairs) was used.

Experimental procedures

20

Plasmid DNA preparation

Total DNA of Human Papilloma Virus type 6 was cloned into vector pSp-64. Plasmid DNA was transferred into E.coli (C-600) and plated onto ampicillin containing LB plates. Single colonies were grown overnight in large culture. Plasmid DNA was isolated according to the method of Birnboim and Doly¹, purified by Sepharose C1-2B column chromatography (Pharmacia) and checked for inserts by restriction-enzyme-analyses. Plasmid DNA concentration was determined by A_{260/280} absorbtion. After ethanol precipitation the DNA was reconstituted in 10mM TRIS/HCl pH 7.2, 0.3mM EDTA to a final concentration of 1 µg/µl (batch# 150894). Subsequently this DNA was sonicated (Soniprep 150., MSE) for 3 times 10 minutes (amplitude 5 microns) on ice.

The size of the resulting DNA fragments was determined by 2% agarose gel electrophoresis and found to be in between 200-400 basepairs (batch# 051094).

5 Plasmid DNA labeling and purification

Plasmid HPV-6 DNA was labeled with BioDadoo-ULS by mixing the following reagents:

plasmid HPV-6 DNA (batch# 051094)	5 μ l (1 μ g/ μ l)
BioDadoo-ULS labeling reagent	8 μ l (1 μ g/ μ l)

10 (batch# BX940830)

Demineralised water (<0.2/ μ S/cm)	37 μ l
--	------------

The 50 μ l reaction mixture was incubated for 15 minutes at 85°C.

Excess of labeling reagent was captured by adding 50 μ l sodium

15 diethyldithiocarbamate (2% solution in demineralised water) and incubating for 30 minutes at room temperature.

Unbound BioDadoo-ULS was removed, using a S300 HR microspin column (Pharmacia), by size exclusion chromatography.

Eluent volume was adapted to 500 μ l giving a 10 ng/ μ l biotin

20 HPV-6 probe concentration (batch# 061094).

Quality control for detection limits

The detection limit of the biotin probe of the invention was determined by direct spot blot and reversed

25 filterhybridization according to the following protocols:

Direct spot blot

HPV-6 probe (batch# 061094) labeled with biotin according to the invention was 10-fold serially diluted into spot buffer

30 comprising 900mM sodium chloride, 90mM sodium citrate and 200 μ g/ml single stranded salmon sperm DNA giving a dilution series varying from 1000-0.1 pg biotin probe per μ l.

1 μ l spots were applied onto nitrocellulose membrane and incubated for 2 hours at 80°C to bind the DNA. The biotin

35 probe was visualized using a streptavidin-alkaline phosphatase conjugate (Sigma) combined with a NBT/BCIP precipitating

substrate solution (Sigma) according to the following protocol:

- Membranes were soaked in TBS solution containing 0.5% tween20 (TBST) for 5 minutes.
- 5 - Membranes were incubated with Strep-AP (3 DEA U/ml) in TBST for 15 minutes at 37°C.
- NC-membranes were washed 3 times 5 minutes in TBS solution followed by a 5 minute wash step in demineralised water.
- Membranes were incubated with NBT/BCIP substrate solution
- 10 for 15 minutes at 37°C, subsequently washed in demineralised water and air dried.

Results

By using this method the detection limit of the biotin DNA
15 probe according to the invention was found to be less than 1 pg.

Reversed filterhybridization

HPV-6 DNA (batch# 051094) was 1 in 10 diluted in 0.1N NaOH,
20 incubated at 100°C for 5 minutes and directly placed on ice for 5 minutes to make DNA single stranded.

A 10-fold serial dilution was made in cold 0.1N NaOH to give a series varying from 10,000-1pg DNA per μ l. 1 μ l spots were
25 applied onto Nylon membrane (Boehringer Mannheim) and air dried.

HPV-6 DNA probe that was labeled with biotin according to the invention was diluted in 5xSSPE 0.5% SDS solution to yield a concentration of 200ng/ml.

This sloution was incubated for 5 minutes at 100°C and placed
30 directly on ice for 5 minutes.

Nylon membranes containing target DNA were soaked in 2x SSC for 5 minutes and subsequently incubated with the single stranded probe solution for 2 hours at 37°C.

Excess of the biotin probe was removed by three changes in 2x
35 SSPE 0.1% SDS for 10 minutes at 37°C followed by a 5 minutes TBST incubation.

The biotin probe of the invention was visualized by performing the same protocol as described in the direct spot blot method.

Results

- 5 By using this procedure the detection limit of the biotin probe according to the invention was found to be less than 10 pg.

Performance in In Situ Hybridization

- 10 The test material consisted of 6 μ m paraffin sections of a HPV-6 positive cervical condyloma mounted on organosilane coated glass slides.

The following protocol was applied (unless otherwise stated steps are at room temperature):

- 15 1 Paraffin sections were dewaxed in 3 changes of xylene and hydrated in graded ethanol.
2 Sections were rinsed in TBST for 5 minutes.
3 Sections were digested in 0.25% pepsin in 0.1N HCl for 30 minutes at 37°C, dehydrated in graded ethanol and air dried.
20 4 10 μ l of probe solution was applied to a section and covered with a coverslip.
Probe solution consisted of biotin HPV-6 probe DNA labeled according to the invention (batch# 061094) in a concentration of 2 ng/ μ l dissolved in hybridization mixture comprising 0.6M
25 NaCl, 0.06M sodium citrate, 35% formamid, 10% dextran sulphate, 2,5x Denhardt's and 10 μ g/ml single stranded salmon sperm DNA.
5 Slides were placed on a hot plate set at 95°C for 5 minutes to denature probe and target DNA simultaneously.
6 Hybridization was performed by placing the slides in a
30 humidified chamber at 37°C for 2 hours.
7 Coverslips were removed and slides were washed in 3 changes of 15mM NaCl, 1.5mM sodium citrate and 5% formamid for 10 minutes at 37°C.
8 Slides were rinsed in TBST.
35 9 Sections were incubated with Streptavidin AP conjugate (3DEA U/ml in TBST) for 15 minutes at 37°C.

10 Slides were washed in TBST (3x) and demineralised water (1x) for 5 minutes.

11 Sections were incubated with NBT/BCIP substrated solution for 15 minutes at 37°C.

5 12 Slides were washed in demineralised water (3x) for 1 minute and sections were mounted in glycerol/gelatin.

Results

10 By using the sections showed blue/purple precipitates at the sites of HPV-6 infected cells and minor background in the remaining tissue.

Conclusions

15 The results demonstrate that DNA labeled according to the invention has good detection limits. The present method is very well suited for research, routine and for industrial production of labeled nucleic acids, as the method is fast and easy to perform, very sensitive, and does not include any enzymatic step, which makes it highly reproducible and fitted
20 for an overall low cost production. The method of the invention offers a useful alternative equaling conventional non-isotopic labeling methods.

General references

- 25 1. Maniatis T., Sambrook J., Fritsch E.F., Molecular Cloning, Second Edition, Cold Spring Harbor Laboratory Press, ISBN 0-8769-309-6.
2. Keller G.H., Manak M.M., DNA probes, Stockton Press, ISBN 0-333-47659-X.

30

Applications

1. The use of Pt-DNA linkers of the invention in the so called LIDIA technique: Linked DNA immuno Assay.

35

The LIDIA technique enables the quantitative analysis of small amounts of DNA (or RNA) e.g. after a PCR amplification of the starting material. The technique is sensitive and

specific, due to the use of specific DNA(RNA) probes in accordance with the invention and easy to perform, because of the quick DNA(RNA) Pt-labeling steps of the invention.

5 Description of the technique:

The technique uses fast Pt labeling compounds of the invention to label DNA(RNA) probes

- 10 This technique is possible with 3 different approaches.
1. Linking DNA probes molecules to a surface by using a Pt compound in accordance with the invention which cross-link DNA molecules irreversibly to plastic, nylon or nitrocellulose. Detection of DNA targets can then be
15 accomplished by using classically labeled DNA/RNA probes.(nick translation or chemical modification, random priming)
 2. Linking a detectable group to the DNA, to render a DNA molecule into a so-called DNA probe. Binding of DNA
20 compounds to a surface can then be accomplished by using classic techniques known to science (covalent linking to specially treated microtiter plates, baking of DNA molecules onto nitrocellulose or binding of DNA molecules to nylon membranes.
 - 25 3. A combination of techniques 1 and 2

Approach 1

- 30 An immobilized DNA probe can be used to catch specific target molecules in a sample by using a hybridization technique. Detection of formed hybrids can be done by using different techniques, e.g. a second labeled DNA probe can be used to hybridize with a different site on the target DNA molecule to form a sandwich hybrid. The label can then be
35 detected by using state of the art immunological detection and colouring techniques.

Approach 2

A volume containing (amplified) detectable DNA(RNA) is directly labeled according to the protocol in accordance with
5 the invention.

Excess label is quenched by adding NaDDTC or Thiourem. This approach distinguishes itself from other techniques by the fact that the target molecule is labeled in contrast to other assay where labeled DNA(RNA) probes are used to detect
10 the target. The quick binding capacity of the Pt-label compound of the invention enables a DNA binding step as a routine step in a diagnostic test procedure (normal binding times are 60 minutes at 65°C).

A second step is performed in a microtiter plate
15 precoated with a target specific probe. Incubation is allowed to the formation of stable "Labelled target" and probe hybrids. The direct labeling of target molecules enables the omission of laborious double hybridization techniques where one probe is used to catch the target and another labeled
20 probe is used to detect the immobilized target.

In this method the probes are covalently linked to the microtiter plate to the surface of the wells. The second incubation step has the character of a liquid hybridisation and therefore can be performed very rapidly. This is one of
25 the main innovative features of this approach to quantitative DNA hybridisation techniques.

Approach 3

30 Both for the immobilization of DNA probes or DNA targets and for the labeling of DNA probes and targets the newly developed Pt system can be used. These two DNA linking techniques can be combined into one assay where both the "catcher" and the "detector" are linked to a second substance
35 (either a detectable group like biotin, digoxigenin or a carrier surface like a plastic stick, microtiter plate or a membrane).

Examples of the technique: the detection of STD related microorganisms in human diagnostics (Chlamydia, Syphilis, AIDS, Herpes, Gonorrhoea, Hep. B,)

5

2. The use of Pt-DNA labels of the invention in combination with test strip procedures and formats. The "DNA Dipstick".

10

The DNA dipstick technique enables the qualitative and semi-quantitative analysis of small amounts of DNA(or RNA) e.g. after a PCR amplification or freely present in samples of body fluids (blood, urine, sweat etc.)

The technique is sensitive and specific, due to the use of specific DNA(RNA) probes and easy to perform because of the quick DNA(RNA) Pt-labeling steps according to the invention.

The universal labeling characteristics of the newly developed Pt label can be used in 3 ways to achieve a bound DNA(RNA) molecule.

20

1. It can be used to attach a detectable marker group to a polynucleotide sequence.
2. It can be used to attach polynucleotide sequences irreversibly to a solid phase (plastic, membranes, latex beads, hydrosols or microtiter plate wells).

25

3. A combination of 1 and 2
ad 1:

In this example there is a twofold approach to the detection of biological analytes in test samples.

30

Firstly a DNA probe can be labeled with the newly developed Pt labeling compound. This labeled probe can then be used to detect preformed hybrids on a membrane formed between the target DNA sequence and a primary probe. It is essential in this method that the primary probe recognizes a different sequence on the target than the secondary Pt labeled probe. In practice, this can be achieved for instance with RNA hybridization where a POLY A probe is used as a primary probe

35

to immobilize all RNA (recognizable by its polyT tails) to a membrane.

The second approach differs slightly in that in this case the target can be labeled in the test sample fluid, because of the fast and very specific Pt labeling characteristics. A procedure like this would comprise a catch of the labeled target with an immobilized specific unlabeled DNA probe on a suitable membrane. Hence a dipstick version for DNA/RNA applications.

10

ad 2:

To immobilize DNA probes or target DNA, a non-labeled Pt compound (that is a Pt compound with 2 free binding sites) can be used to act as a bridge between DNA and the surface of carriers (plastic, membrane, microtiter plates etc.)

15

It greatly enhances the usability of DNA sequences as catcher molecules in diagnostic assays, since there are little substances known to science that bind readily DNA in a spontaneous way. Introducing this Pt bridge molecule a wide field of new applications for the DNA technology has come within reach.

20

ad 3: a combination of example 1 and 2

General: the use of the Pt compound of the invention in latex or hydrosol assays is particularly interesting. The compound enables the coupling of DNA molecules to small particles. The DNA molecules can be hybridized to target material. A positive reaction is visualized by an agglutination of the particles, due to crosslinking of the DNA hybrid particle compounds.

30

A test like this can be made quantative, the rate of agglutination can be tuned and measured at a specific wavelength. Especially gold particles have the intrinsic characteristic that a shift in optimal wavelength occurs after agglutination.

35

3. Detection of Platinated DNA probes of the invention with the silver-enhancement technique.

Platinated DNA/RNA probe can be employed in hybridisation methods to detect DNA/RNA sequences in sample material. The introduction of a platinum compound at the site of the target enables the deposition of Ag molecules in a chemical reaction especially designed to reduce ionic silver to metallic silver. At the site of a Pt nucleus a decomposition of metallic silver(black) occurs due to the catalytic effect of the Pt nucleus.


Ionic silver is reduced by a reducing agent (e.g. Na-borohydrid, Hydrochinon) in solution. In a constant ratio the amount of silver deposited on the Pt is proportional to the length of the enhancement incubation. Visualisation of a non-visible Pt nucleus can be accomplished by the empirical observation of the appearance of a black spot in the test sample.

The black spots indicate the site of specific probes binding and thus the site of specific target location. The technique enables a quick and easy diagnostic procedure for the detection of various microorganisms and gene translocations/abnormalities.

CLAIMS

1. A method for labeling nucleotides comprising the steps of:

- reacting a reactive moiety of a linker of the formula



5

wherein X represents any stabilizing bridge and wherein A and B represent the same or different reactive moieties, with an electron donating moiety of a spacer, which spacer comprises a chain having at least four atoms and at least one heteroatom in the chain, which spacer further comprises said electron donating moiety at one end of the chain and a reactive moiety at the other end of the chain;

10

- reacting the reactive moiety of said spacer with a label;
- 15 - reacting the other reactive moiety of said linker with a nucleotide.

2. A method according to claim 1 wherein the label is biotin, avidin, streptavidin, digoxigenin or a functional equivalent thereof.

20

3. A method according to claim 1 or 2 wherein the nucleotide is adenine, thymidine, cytosine, guanine or uridine or a derivative thereof.


25

4. A method according to any of the preceding claims wherein the electron donating moiety is an amine or a thiolate anion.


5. A method according to claim 4 wherein the amine is an aromatic amine.

30

6. A method according to any of the preceding claims wherein at least one heteroatom is an oxygen atom.
- 5 7. A method according to any of the preceding claims wherein the spacer comprises no more than 20 carbon atoms in the chain and wherein the chain is essentially non-branched.
8. A method according to claim 7 wherein the spacer is 1,8-
10 diamino-3,6-dioxaoctane.
9. A method according to claim 7 wherein the spacer is an oligolysine or a polylysine.
- 15 10. A method according to claims 1-4 wherein the linker is reacted with a labeling moiety comprising a label and a spacer, which labeling moiety has the formula



20 or the formula



wherein X represents any stabilizing bridge, Z and Z' represent a non-leaving ligand and n is an integer of from 2
 5 to 10.

11. A method according to claim 10, wherein Z and/or Z' represent an NH_3 , NH_2R , NHR_2 , or NR_3 group, wherein R represents an alkyl group having from 1 to 6 carbon atoms.
 10

12. A method according to any of the preceding claims, wherein X represents an aliphatic diamine.


13. A method according to claim 12 wherein X represents an aliphatic diamine having 2-6 carbon atoms.
 15

14. A method according to claim 13 wherein X is an ethylene diamine group.

20 15. A method according to claims 12-14 wherein one or both of the nitrogen atoms of the aliphatic diamine are shielded.

16. A method according to claim 15 wherein one or both of the nitrogen atoms of the aliphatic diamine are substituted with
 25 an alkyl group of from 1 to 6 carbon atoms.

17. A method according to claim 16 wherein one or both of the nitrogen atoms of the aliphatic diamine are substituted with one or two methyl groups.
- 5 18. A method according to any of the preceding claims wherein A and/or B are selected from the group consisting of NO_3^- , SO_3^- , Cl^- , I^- , or other halogens.
19. A method according to any of the preceding claims wherein
10 A and B are the same.
20. A labeled nucleotide obtainable by a method according to any of the preceding claims.
- 15 21. A labeled nucleotide according to claim 20 wherein the spacer is an oligolysine or a polylysine.
22. A labeling substance having the formula



20

- wherein X represents any stabilizing bridge, A represents a reactive moiety and the labeling moiety comprises a label and a spacer, which spacer comprises a chain having at least four
25 atoms and at least one heteroatom in the chain, which spacer further comprises an electron donating moiety at the end of the spacer distal from the label.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/NL 97/00559

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07F15/00 C07H21/00 C07H23/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 01699 A (ACADEMISCH MEDISCH CENTRUM ;UNIV LEIDEN (NL)) 6 February 1992 cited in the application see the whole document ---	1,22
P,X	WO 96 35696 A (KREATECH BIOTECH BV ;HOETHOFF HENDRIK JAN (NL); REEDIJK JAN (NL);) 14 November 1996 see the whole document ---	1,20,22
A	GB 2 148 891 A (NIPPON KAYAKU KK) 5 June 1985 see the whole document ---	1,20,22
A	US 4 207 416 A (HOESCHELE JAMES D) 10 June 1980 see the whole document ---	1,20,22
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 November 1997

Date of mailing of the international search report

05/12/1997

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Moreno, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 97/00559

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 282 672 A (NIPPON KAYAKU KK) 21 September 1988 see the whole document ---	1,20,22
A	EP 0 386 243 A (SAGAMI CHEM RES ;CHISSO CORP (JP)) 12 September 1990 see the whole document -----	1,20,22

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. l. Application No

PCT/NL 97/00559

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9201699 A	06-02-92	NL 9001639 A AT 145403 T AU 8286391 A DE 69123251 D DE 69123251 T EP 0539466 A ES 2097213 T US 5580990 A	17-02-92 15-12-96 18-02-92 02-01-97 28-05-97 05-05-93 01-04-97 03-12-96
WO 9635696 A	14-11-96	AU 5704096 A	29-11-96
GB 2148891 A	05-06-85	JP 1725766 C JP 4012277 B JP 60087295 A BE 900842 A DE 3438440 A DK 499584 A FR 2553777 A NL 8403087 A SE 8405188 A US 4607114 A	19-01-93 04-03-92 16-05-85 18-04-85 02-05-85 20-04-85 26-04-85 17-05-85 20-04-85 19-08-86
US 4207416 A	10-06-80	AU 8455375 A CA 1034120 A CH 617587 A DE 2539179 A FR 2283692 A GB 1526210 A JP 51054916 A SE 424870 B SE 7509769 A US 4080324 A	10-03-77 04-07-78 13-06-80 18-03-76 02-04-76 27-09-78 14-05-76 16-08-82 08-03-76 21-03-78
EP 0282672 A	21-09-88	JP 63203692 A CA 1292749 A US 4861905 A US 4987246 A US 5128493 A	23-08-88 03-12-91 29-08-89 22-01-91 07-07-92
EP 0386243 A	12-09-90	JP 2000294 A	05-01-90

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 97/00559

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0386243 A		JP 2033479 C	19-03-96
		JP 7062023 B	05-07-95
		JP 1249791 A	05-10-89
		WO 8904317 A	18-05-89
<hr/>			